



Choudhury, S., Lightman, S., & Meeran, K. (2019). Improving glucocorticoid replacement profiles in adrenal insufficiency. *Clinical Endocrinology*, 91(3), 367-371. <https://doi.org/10.1111/cen.13999>

Peer reviewed version

Link to published version (if available):  
[10.1111/cen.13999](https://doi.org/10.1111/cen.13999)

[Link to publication record in Explore Bristol Research](#)  
PDF-document

This is the author accepted manuscript (AAM). The final published version (version of record) is available online via Wiley at <https://onlinelibrary.wiley.com/doi/full/10.1111/cen.13999> . Please refer to any applicable terms of use of the publisher.

## University of Bristol - Explore Bristol Research

### General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:  
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>



## Improving glucocorticoid replacement profiles in adrenal insufficiency

Journal:	<i>Clinical Endocrinology</i>
Manuscript ID	CEN-2019-000070.R3
Manuscript Type:	6 Requested Review
Date Submitted by the Author:	n/a
Complete List of Authors:	Choudhury, Sirazum; Imperial College London, Section of Investigative Medicine; Imperial College Healthcare NHS Trust, Department of Clinical Biochemistry Lightman, Stafford; University of Bristol, Henry Wellcome Laboratories for Integrative Neuroscience and Endocrinology Meeran, Karim; Imperial College London, Section of Investigative Medicine; Imperial College Healthcare NHS Trust, Department of Endocrinology
Key Words:	Adrenal insufficiency < Conditions: < Adrenal, Glucocorticoids < Hormones/related: < Adrenal, Cortisol < Hormones/related: < Adrenal, Prednisolone < Investigations & Rx: < Adrenal

**Improving glucocorticoid replacement profiles in adrenal insufficiency**

**Glucocorticoid replacement in AI**

Sirazum Choudhury MBBS<sup>1</sup>, Stafford Lightman FRS<sup>2</sup> and Karim Meeran FRCP FRCPath<sup>1</sup>

<sup>1</sup>Department of Investigative Medicine, Division of Diabetes, Endocrinology and Metabolism, Imperial College London, Commonwealth Building, London, UK

<sup>2</sup>Henry Wellcome Laboratories for Integrative Neuroscience and Endocrinology, University of Bristol, Dept of Translational Health Science, Bristol, UK

**Corresponding author:**

Professor Karim Meeran

Department of Investigative Medicine, 6<sup>th</sup> Floor Commonwealth Building, Imperial College London, Du Cane Road London, W12 0NN

Email: k.meeran@imperial.ac.uk

Telephone: 020 3311 1065      Fax: 020 3311 1862

**Acknowledgements**

Nil

**Conflict of Interest:**

None of the authors declare a conflict of interest.

**Summary:**

There is an increased mortality associated with adrenal insufficiency despite glucocorticoid replacement therapy with a standardised mortality ratio greater than two. The cause of the increased mortality is yet to be definitively elucidated, but may be due to excess steroid exposure, or replacement regimens that are uncoupled from the normal physiological cortisol profile. Cortisol secretion follows an ultradian pattern which is not possible to reproduce using oral replacement. With the advent of new pumps, it is now possible to mimic the pulsatility of the adrenal glands. Whilst the cognitive and emotional benefits of reproducing the ultradian rhythm is known, the presence of long-term benefits are not yet clear.

There is a dearth of evidence and high-quality studies to underline our current understanding of the pathophysiology of adrenal insufficiency and replacement therapy. There is a particular lack of research comparing objective outcomes between patients receiving hydrocortisone replacement (either standard therapy or new sustained release preparations), prednisolone replacement, and ultradian pumps. Direct comparative studies are now warranted to understand the optimal approach.

**Keywords:**

Adrenal insufficiency, hypocortisolism, glucocorticoids, cortisol, hydrocortisone, prednisolone.

**Main Text:**

There is no doubt that steroid replacement with glucocorticoids saves lives in patients with adrenal insufficiency, a condition with an associated mortality of 100% before the development of glucocorticoids<sup>1</sup>. Excess glucocorticoid exposure, as evidenced by patients with Cushing's syndrome, can also be fatal. Remarkably however, there is no agreement on how we can optimise replacement therapy for either primary or secondary adrenal insufficiency. Several options result in excess glucocorticoid exposure, which may reduce the risk of an Addisonian crisis, but only at the cost of a reduced natural killer cell function<sup>2</sup>, an increased cardiovascular disease burden<sup>3,4</sup>, and increased mortality<sup>5</sup>. Whether the increased mortality is the result of excessive exposure to glucocorticoid, or inappropriate timing and rhythm of replacement remains unclear.

The importance of circadian rhythmicity has been clear for a long time and life forms from primitive cyanobacteria, through all phyla to man, have adapted to this with mechanisms to predict the changing environments of the 24-hour day. Uncoupling glucocorticoid dose timing from the normal circadian cortisol rhythm is harmful<sup>6,7</sup> and human cells have developed circadian clocks that are, to varying extents, kept in synchrony by the HPA axis<sup>8</sup>. Cortisol levels rise at around 4am, under the influence of the central circadian pacemaker in the hypothalamic suprachiasmatic nucleus. This activates or inhibits a massive variety of genes in all tissues of the body, in anticipation of the body's needs at awakening and can also alter the endogenous clock gene "setting" of cells<sup>9</sup>. Misalignment between cortisol levels and clock time is one of the factors contributing to the adverse health outcomes seen in shift workers and problems associated with jet lag<sup>10</sup>.

***Oral preparations of glucocorticoids***

The excess replacement that frequently occurs with twice or thrice daily hydrocortisone is associated with an increase in cardiovascular deaths in these patients, especially with escalating doses<sup>4</sup> and the late peaks in cortisol might be harmful<sup>11</sup>. A delayed release preparation (Chronocort) taken last thing at night<sup>12</sup> has the theoretical advantage of mimicking the early morning 4am rise in cortisol, which cannot be achieved with other preparations. Chronocort has already shown potential to suppress morning and 24-hour androstenedione levels in patients with congenital adrenal hyperplasia during phase II trials<sup>13</sup>, but more studies are needed to determine the clinical importance of this anticipatory rise in cortisol.

Another strategy available in some countries is the use of dual release hydrocortisone (Plenadren) once daily first thing in the morning. This sustained release preparation fails to replicate the pre-awakening rise in cortisol and we know very little about what happens to blood or tissue levels of cortisol overnight. It relies on continuous absorption of hydrocortisone from the gut which can be variable, a problem that must be guarded against in patients with infectious diarrhoea, when glucocorticoids are crucial<sup>14</sup>. The manufacturers state that Plenadren is commonly associated (> 1 in 10 cases) with gastrointestinal side effects<sup>15</sup>.

Building on the early comparative findings that Plenadren may lower blood pressure, weight and improve HbA1c compared to multiple daily cortisone acetate or hydrocortisone<sup>16</sup>, the DREAM study undertook a similar comparison<sup>17</sup>. Although the primary outcome focussed on weight change, the DREAM study also looked at the effect of Plenadren on immune profiles. Recruiting 89 patients on multiple daily cortisone acetate or hydrocortisone, the investigators switched 46 patients to

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Plenadren and asked 43 to continue on the thrice daily immediate release preparations. A third group of 25 healthy volunteers were also recruited and data collected over a 24-week follow up period. In addition to a significant corrected weight loss of 4Kg in the switch group, the investigators reported significantly higher classical (CD14+CD16-) monocytes, with lower non-classical (CD14-CD16+) monocytes and CD16+ mature natural killer cells in patients taking thrice-daily hydrocortisone at baseline versus the healthy volunteers. Converting patients to once daily Plenadren was associated with improved numbers of the classical and non-classical monocytes and the mature natural killer cells, with concordant decreases in the soluble CD16 immune marker (an indicator of natural killer cell immaturity) and reported rates of infection as measured by a questionnaire<sup>17</sup>. These findings suggest that Plenadren may be less immunosuppressive than multiple daily dosing of glucocorticoids and that leucocytes may hold the key to assessing the adequacy of an individual's glucocorticoid replacement therapy. Because a majority of the patients were on cortisone acetate, it is not clear whether the variable metabolism of cortisone to cortisol might be a factor.

A DREAM ancillary study also investigated the effect of the switch to Plenadren on the RNA expression of clock genes. The analysis found that 19 of 68 different clock genes were differently expressed at baseline in all patients taking thrice daily hydrocortisone compared to the healthy volunteer control group<sup>18</sup>. In the switch group, the expression of 18 of the 19 affected genes were found to normalised to the levels seen within the healthy volunteers over 12 weeks. Clock genes are involved in auto-regulated feedback cycles thought to be responsible for conveying clock time to individual tissues and cells. The oscillations of clock genes regulate and control the transcription of proteins and can inform tissue sensitivity to glucocorticoids depending on the time of day, through acetylation of lysine residues on the glucocorticoid receptor<sup>6,19</sup>. Although dissociation of peripheral clock genes and the central suprachiasmatic clock is associated with disease states, the functional significance is not understood<sup>19</sup>. The physiological improvements demonstrated by Plenadren have been attributed to its ability to better mirror the natural circadian cortisol profile, compared to multiple daily hydrocortisone or cortisone acetate although this association might simply be due to the lower steroid exposure seen with Plenadren, with data indicating a reduction of approximately 20% in area-under-the curve exposure<sup>16</sup>. A majority of the patients were on cortisone acetate, which depends on hepatic activation by 11 $\beta$  hydroxysteroid dehydrogenase (11 $\beta$ HSD1) to cortisol<sup>20</sup>. It was assumed that 1mg of cortisone acetate was equivalent to 0.8mg of hydrocortisone<sup>17</sup> but, there is a large (21% to 95%) variability in bioavailability of cortisone acetate<sup>21</sup> as well as tissue specific expression of 11 $\beta$ HSD1.

Prednisolone<sup>22</sup> has a similar plasma profile to Plenadren and since this is taken on waking, the 4am anticipatory rise in glucocorticoid does not occur. At a molecular level we know that prednisolone has prolonged binding to the glucocorticoid receptor compared to hydrocortisone, which would be expected to result in different transcriptional responses in glucocorticoid responsive tissues and prolonged effect<sup>23</sup>. Whether this theoretical advantage is clinically useful and results in reduced risk of adrenal crisis is unknown. The rate of adrenal crises has grown with the increased prescription of short-acting glucocorticoids such as hydrocortisone, although a causal link needs to be confirmed<sup>24</sup>. Further data from a prospective observational study in Germany recorded the frequency of crises experienced by patients on different treatment regimens, and showed that the proportion of patients on prednisolone experiencing a crisis was not greater than expected<sup>25</sup>. It is however clear that previously used doses have been excessive and 3mg to 4mg once daily is sufficient replacement in most patients<sup>26,27</sup>. Currently there are two NIHR adopted studies actively recruiting, comparing low-dose prednisolone with standard regimens of hydrocortisone<sup>28,29</sup>. Low dose prednisolone has been shown to be superior in reducing androgens, 17-hydroxy progesterone and improving growth velocity in patients with congenital adrenal hyperplasia compared to thrice daily hydrocortisone<sup>30</sup>.

### *Subcutaneous glucocorticoid infusions*

Infusion pumps have been used to mimic the circadian rise in morning cortisol. In addition to normalising morning ACTH levels in patients with primary adrenal failure, improvements in physical functioning and vitality domains of the SF-36 have been noted<sup>31,32</sup>. There are however no further anthropometric or objective benefits, and in double-blind placebo-controlled conditions when compared to oral hydrocortisone treatment, the subjective health outcome benefits are no longer detectable<sup>33</sup>. These studies were performed without any pulsatile or ultradian patterns.

Like many other hormones cortisol is secreted in a pulsatile manner<sup>34,35</sup>. This results in a circadian rhythm that is made up of underlying oscillations which reach maximum amplitude at the time of the circadian peak. In animal models these pulses have been shown to be very important for normal physiological and behavioural activity<sup>36,37</sup>. Although there is relatively little variation of circadian rhythm between subjects, there is much more intra-individual variation in the daily pattern of ultradian pulses<sup>34</sup>. Since the frequency of ultradian pulses simply depends on fixed delays in the feedforward-feedback relationship between the pituitary corticotropes and the adrenal fasciculata cells<sup>38,39</sup> the circadian change in cortisol is determined by changes in pulse amplitude rather than pulse frequency. This provides us with a much easier opportunity to use pumps to replicate physiological pulses, as demonstrated by Russell and colleagues<sup>40</sup>.

It is not yet clear to what extent it is important to replicate these rhythms to provide optimal treatment for patients with adrenal failure. Although we know that stress increases the size of ultradian pulses<sup>41,42</sup>, there have been no long term studies on the modulation of ultradian rhythmicity in normal daily living. Physiological studies have shown that it is the presence of both peaks and troughs of cortisol that is important for optimal regulation of behaviour and of glucocorticoid and mineralocorticoid receptors in animals<sup>43,44</sup>. We are now testing this in new psychological paradigms in man.

From a patient's perspective infusion pumps are less convenient than tablets. However, the technological advances of new miniature nanopumps either external or with subcutaneous reservoirs will make pumps less intrusive. Kalafatakis and colleagues recently undertook a double-blind randomised crossover study, in which they recruited 15 young males and implemented a "block and replace" approach to cortisol using three distinct regimens to investigate the comparative effects of pulsatility on cognition, emotion and sleep<sup>45</sup>. Participants had their endogenous cortisol secretion blocked using metyrapone and received 20mg of hydrocortisone in three modalities: 1- a subcutaneous pulsatile mode to replicate the ultradian profile; 2- a subcutaneous cutaneous infusion; 3- oral thrice daily replacement, each for a five-day period. The order in which the modalities were administered was randomised.

The study demonstrated that pulsatility directly affects neurological physiology in the non-stressed state, independent of the cumulative glucocorticoid exposure. Specifically, subcutaneous ultradian pulsatile infusions promoted better quality of sleep and improved performance of working memory compared to the continuous infusions, as assessed by a sleep questionnaire and functional tests respectively.

These different patterns of cortisol oscillations also had differential effects on the fMRI and functional connectivity of the brain regions which process emotional responses (amygdala, dorsal striatum, insula and orbitofrontal cortex) – with parallel changes detected in attentional bias to, and recognition accuracy of emotional cues<sup>45</sup>.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

These data imply that oscillating levels of cortisol are important for physiological regulation of glucocorticoid receptors and post receptor signalling in the brain and for the regulation of mood in man. Whether differences in endogenous patterns of pulsatility in different individuals alters these functions is unknown. The importance of these pulses in longer term replacement therapy in patients with Addison’s disease will be clearer once the Medical Research Council PULSES trial is complete (MR/R010919/1).

Neither immediate release hydrocortisone nor any other oral glucocorticoid will ever replace normal ultradian secretion which remains highly variable across individuals.

**Conclusion**

Long term patient health and cognitive and emotional state as well as patient convenience and compliance are crucial and a balance between a once daily oral dosing, more frequent oral dosing and the requirement of a pump needs to be considered. Currently there are no good studies comparing low dose prednisolone, dual release hydrocortisone (Plenadren, Duocort), delayed release hydrocortisone (Chronocort) and miniature ultradian pumps with three times daily hydrocortisone, using currently understood lower and biologically equivalent doses. We hope that well controlled, relatively long-term double-blind studies will soon be available to allow a rational approach to optimal therapy on an individual patient basis.

**References:**

1. DUNLOP D. Eighty-six cases of addison's disease. *Br Med J*. 1963;2(5362):887-891.

2. Bancos I, Hazeldine J, Chortis V, et al. Primary adrenal insufficiency is associated with impaired natural killer cell function: A potential link to increased mortality. *Eur J Endocrinol*. 2017;176(4):471-480.

3. Behan LA, Carmody D, Rogers B, et al. Low-dose hydrocortisone replacement is associated with improved arterial stiffness index and blood pressure dynamics in severely adrenocorticotrophin-deficient hypopituitary male patients. *Eur J Endocrinol*. 2016;174(6):791-799.

4. Sherlock M, Reulen RC, Alonso AA, et al. ACTH deficiency, higher doses of hydrocortisone replacement, and radiotherapy are independent predictors of mortality in patients with acromegaly. *J Clin Endocrinol Metab*. 2009;94(11):4216-4223.

- 1  
2  
3 5. Hammarstrand C, Ragnarsson O, Hallen T, et al. Higher glucocorticoid replacement doses are  
4 associated with increased mortality in patients with pituitary adenoma. *Eur J Endocrinol*.  
5  
6 2017;177(3):251-256.  
7  
8  
9
- 10  
11 6. Nicolaides NC, Charmandari E, Kino T, Chrousos GP. Stress-related and circadian secretion and  
12 target tissue actions of glucocorticoids: Impact on health. *Front Endocrinol (Lausanne)*. 2017;8:70.  
13  
14  
15
- 16 7. Charmandari E, Nicolaides NC, Chrousos GP. Adrenal insufficiency. *Lancet*. 2014;383(9935):2152-  
17 2167.  
18  
19  
20
- 21 8. Charmandari E, Chrousos GP, Lambrou GI, et al. Peripheral CLOCK regulates target-tissue  
22 glucocorticoid receptor transcriptional activity in a circadian fashion in man. *PLoS One*.  
23  
24 2011;6(9):e25612.  
25  
26  
27
- 28 9. Cuesta M, Cermakian N, Boivin DB. Glucocorticoids entrain molecular clock components in human  
29 peripheral cells. *FASEB J*. 2015;29(4):1360-1370.  
30  
31  
32
- 33 10. Monk TH, Buysse DJ. Exposure to shift work as a risk factor for diabetes. *J Biol Rhythms*.  
34  
35 2013;28(5):356-359.  
36  
37  
38
- 39 11. Muller L, Quinkler M. Adrenal disease: Imitating the cortisol profile improves the immune  
40 system. *Nat Rev Endocrinol*. 2018;14(3):137-139.  
41  
42  
43
- 44 12. Whitaker M, Debono M, Huatan H, Merke D, Arlt W, Ross RJ. An oral multiparticulate, modified-  
45 release, hydrocortisone replacement therapy that provides physiological cortisol exposure. *Clin*  
46  
47 *Endocrinol (Oxf)*. 2014;80(4):554-561.  
48  
49  
50
- 51 13. Mallappa A, Sinaii N, Kumar P, et al. A phase 2 study of chronocort, a modified-release  
52 formulation of hydrocortisone, in the treatment of adults with classic congenital adrenal hyperplasia.  
53  
54  
55  
56  
57  
58  
59  
60 *J Clin Endocrinol Metab*. 2015;100(3):1137-1145.



- 1  
2  
3 14. Allolio B. Extensive expertise in endocrinology. adrenal crisis. *Eur J Endocrinol*. 2015;172(3):R115-  
4  
5 24.  
6  
7  
8  
9 15. Shire Pharmaceuticals. Summary of product characteristics.  
10  
11 [https://www.ema.europa.eu/documents/product-information/plenadren-epar-product-](https://www.ema.europa.eu/documents/product-information/plenadren-epar-product-information_en.pdf)  
12 [information\\_en.pdf](https://www.ema.europa.eu/documents/product-information/plenadren-epar-product-information_en.pdf).  
13  
14  
15  
16  
17 16. Johannsson G, Nilsson AG, Bergthorsdottir R, et al. Improved cortisol exposure-time profile and  
18 outcome in patients with adrenal insufficiency: A prospective randomized trial of a novel  
19 hydrocortisone dual-release formulation. *J Clin Endocrinol Metab*. 2012;97(2):473-481.  
20  
21  
22  
23  
24 17. Isidori AM, Venneri MA, Graziadio C, et al. Effect of once-daily, modified-release hydrocortisone  
25 versus standard glucocorticoid therapy on metabolism and innate immunity in patients with adrenal  
26 insufficiency (DREAM): A single-blind, randomised controlled trial. *Lancet Diabetes Endocrinol*. 2017.  
27  
28  
29  
30  
31  
32 18. Venneri MA, Hasenmajer V, Fiore D, et al. Circadian rhythm of glucocorticoid administration  
33 entrains clock genes in immune cells: A DREAM trial ancillary study. *J Clin Endocrinol Metab*.  
34 2018;103(8):2998-3009.  
35  
36  
37  
38  
39  
40 19. Oster H, Challet E, Ott V, et al. The functional and clinical significance of the 24-hour rhythm of  
41 circulating glucocorticoids. *Endocr Rev*. 2017;38(1):3-45.  
42  
43  
44  
45  
46 20. Fariss BL, Hane S, Shinsako J, Forsham PH. Comparison of absorption of cortisone acetate and  
47 hydrocortisone hemisuccinate. *J Clin Endocrinol Metab*. 1978;47(5):1137-1140.  
48  
49  
50  
51 21. Heazelwood VJ, Galligan JP, Cannell GR, Bochner F, Mortimer RH. Plasma cortisol delivery from  
52 oral cortisol and cortisone acetate: Relative bioavailability. *Br J Clin Pharmacol*. 1984;17(1):55-59.  
53  
54  
55  
56  
57  
58  
59  
60

22. Williams EL, Choudhury S, Tan T, Meeran K. Prednisolone replacement therapy mimics the circadian rhythm more closely than other glucocorticoids. *The Journal of Applied Laboratory Medicine*. 2016;1(2):152-161.
23. Stavreva DA, Wiench M, John S, et al. Ultradian hormone stimulation induces glucocorticoid receptor-mediated pulses of gene transcription. *Nat Cell Biol*. 2009;11(9):1093-1102.
24. Rushworth RL, Torpy DJ. Adrenal insufficiency in australia: Is it possible that the use of lower dose, short-acting glucocorticoids has increased the risk of adrenal crises? *Horm Metab Res*. 2015;47(6):427-432.
25. Hahner S, Spinnler C, Fassnacht M, et al. High incidence of adrenal crisis in educated patients with chronic adrenal insufficiency: A prospective study. *J Clin Endocrinol Metab*. 2015;100(2):407-416.
26. Smith DJF, Prabhudev H, Choudhury S, Meeran K. Prednisolone has the same cardiovascular risk profile as hydrocortisone in glucocorticoid replacement. *Endocr Connect*. 2017;6(8):766-772.
27. Choudhury S, Machenahalli P, Tan T, Meeran K. Inadvertent treatment of hypoadrenalism with prednisolone in pemphigus: A case report. *Clin Case Rep*. 2019;0(0).
28. Imperial College London. Safety and efficacy of prednisolone in adrenal insufficiency disease (PRED-AID study). ISRCTN Registry Web site. <http://www.isrctn.com/ISRCTN41325341>. Published 01/28/2019. Updated 2019. Accessed 04/19, 2019.
29. Imperial College London. **Hydrocortisone vs prednisolone in AI (HYPER-AID)**. Clinicaltrials.gov Web site. <https://clinicaltrials.gov/ct2/show/NCT03608943>. Published 08/01/2018. Updated 2019. Accessed 04/19, 2019.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

30. Caldato MC, Fernandes VT, Kater CE. One-year clinical evaluation of single morning dose prednisolone therapy for 21-hydroxylase deficiency. *Arq Bras Endocrinol Metabol.* 2004;48(5):705-712.

31. Lovas K, Husebye ES. Continuous subcutaneous hydrocortisone infusion in addison's disease. *Eur J Endocrinol.* 2007;157(1):109-112.

32. Oksnes M, Bjornsdottir S, Isaksson M, et al. Continuous subcutaneous hydrocortisone infusion versus oral hydrocortisone replacement for treatment of addison's disease: A randomized clinical trial. *J Clin Endocrinol Metab.* 2014;99(5):1665-1674.

33. Gagliardi L, Nenke MA, Thynne TR, et al. Continuous subcutaneous hydrocortisone infusion therapy in addison's disease: A randomized, placebo-controlled clinical trial. *J Clin Endocrinol Metab.* 2014;99(11):4149-4157.

34. Lightman S, Terry JR. The importance of dynamic signalling for endocrine regulation and drug development: Relevance for glucocorticoid hormones. *Lancet Diabetes Endocrinol.* 2014;2(7):593-599.

35. Faghih RT, Dahleh MA, Adler GK, Klerman EB, Brown EN. Deconvolution of serum cortisol levels by using compressed sensing. *PLoS One.* 2014;9(1):e85204.

36. Lightman SL, Conway-Campbell BL. The crucial role of pulsatile activity of the HPA axis for continuous dynamic equilibration. *Nat Rev Neurosci.* 2010;11(10):710-718.

37. Spiga F, Walker JJ, Terry JR, Lightman SL. HPA axis-rhythms. *Compr Physiol.* 2014;4(3):1273-1298.

38. Walker JJ, Terry JR, Lightman SL. Origin of ultradian pulsatility in the hypothalamic-pituitary-adrenal axis. *Proc Biol Sci.* 2010;277(1688):1627-1633.

39. Walker JJ, Spiga F, Waite E, et al. The origin of glucocorticoid hormone oscillations. *PLoS Biol.* 2012;10(6):e1001341.
40. Russell GM, Durant C, Ataya A, et al. Subcutaneous pulsatile glucocorticoid replacement therapy. *Clin Endocrinol (Oxf).* 2014;81(2):289-293.
41. Henley DE, Russell GM, Douthwaite JA, et al. Hypothalamic-pituitary-adrenal axis activation in obstructive sleep apnea: The effect of continuous positive airway pressure therapy. *J Clin Endocrinol Metab.* 2009;94(11):4234-4242.
42. Gibbison B, Spiga F, Walker JJ, et al. Dynamic pituitary-adrenal interactions in response to cardiac surgery. *Crit Care Med.* 2015;43(4):791-800.
43. Sarabdjitsingh RA, Conway-Campbell BL, Leggett JD, et al. Stress responsiveness varies over the ultradian glucocorticoid cycle in a brain-region-specific manner. *Endocrinology.* 2010;151(11):5369-5379.
44. George CL, Birnie MT, Flynn BP, Kershaw YM, Lightman SL, Conway-Campbell BL. Ultradian glucocorticoid exposure directs gene-dependent and tissue-specific mRNA expression patterns in vivo. *Mol Cell Endocrinol.* 2017;439:46-53.
45. Kalafatakis K, Russell GM, Harmer CJ, et al. Ultradian rhythmicity of plasma cortisol is necessary for normal emotional and cognitive responses in man. *Proc Natl Acad Sci U S A.* 2018;115(17):E4091-E4100.

**Improving glucocorticoid replacement profiles in adrenal insufficiency**

**Glucocorticoid replacement in AI**

Sirazum Choudhury MBBS<sup>1</sup>, Stafford Lightman FRS<sup>2</sup> and Karim Meeran FRCP FRCPath<sup>1</sup>

<sup>1</sup>Department of Investigative Medicine, Division of Diabetes, Endocrinology and Metabolism, Imperial College London, Commonwealth Building, London, UK

<sup>2</sup>Henry Wellcome Laboratories for Integrative Neuroscience and Endocrinology, University of Bristol, Dept of Translational Health Science, Bristol, UK

**Corresponding author:**

Professor Karim Meeran

Department of Investigative Medicine, 6<sup>th</sup> Floor Commonwealth Building, Imperial College London, Du Cane Road London, W12 0NN

Email: k.meeran@imperial.ac.uk

Telephone: 020 3311 1065      Fax: 020 3311 1862

**Acknowledgements**

Nil

**Conflict of Interest:**

None of the authors declare a conflict of interest.

**Summary:**

There is an increased mortality associated with adrenal insufficiency despite glucocorticoid replacement therapy with a standardised mortality ratio greater than two. The cause of the increased mortality is yet to be definitively elucidated, but may be due to excess steroid exposure, or replacement regimens that are uncoupled from the normal physiological cortisol profile. Cortisol secretion follows an ultradian pattern which is not possible to reproduce using oral replacement. With the advent of new pumps, it is now possible to mimic the pulsatility of the adrenal glands. Whilst the cognitive and emotional benefits of reproducing the ultradian rhythm is known, the presence of long-term benefits are not yet clear.

There is a dearth of evidence and high-quality studies to underline our current understanding of the pathophysiology of adrenal insufficiency and replacement therapy. There is a particular lack of research comparing objective outcomes between patients receiving hydrocortisone replacement (either standard therapy or new sustained release preparations), prednisolone replacement, and ultradian pumps. Direct comparative studies are now warranted to understand the optimal approach.

**Keywords:**

Adrenal insufficiency, hypocortisolism, glucocorticoids, cortisol, hydrocortisone, prednisolone.

**Main Text:**

There is no doubt that steroid replacement with glucocorticoids saves lives in patients with adrenal insufficiency, a condition with an associated mortality of 100% before the development of glucocorticoids<sup>1</sup>. Excess glucocorticoid exposure, as evidenced by patients with Cushing's syndrome, can also be fatal. Remarkably however, there is no agreement on how we can optimise replacement therapy for either primary or secondary adrenal insufficiency. Several options result in excess glucocorticoid exposure, which may reduce the risk of an Addisonian crisis, but only at the cost of a reduced natural killer cell function<sup>2</sup>, an increased cardiovascular disease burden<sup>3,4</sup>, and increased mortality<sup>5</sup>. Whether the increased mortality is the result of excessive exposure to glucocorticoid, or inappropriate timing and rhythm of replacement remains unclear.

The importance of circadian rhythmicity has been clear for a long time and life forms from primitive cyanobacteria, through all phyla to man, have adapted to this with mechanisms to predict the changing environments of the 24-hour day. Uncoupling glucocorticoid dose timing from the normal circadian cortisol rhythm is harmful<sup>6,7</sup> and human cells have developed circadian clocks that are, to varying extents, kept in synchrony by the HPA axis<sup>8</sup>. Cortisol levels rise at around 4am, under the influence of the central circadian pacemaker in the hypothalamic suprachiasmatic nucleus. This activates or inhibits a massive variety of genes in all tissues of the body, in anticipation of the body's needs at awakening and can also alter the endogenous clock gene "setting" of cells<sup>9</sup>. Misalignment between cortisol levels and clock time is one of the factors contributing to the adverse health outcomes seen in shift workers and problems associated with jet lag<sup>10</sup>.

*Oral preparations of glucocorticoids*

The excess replacement that frequently occurs with twice or thrice daily hydrocortisone is associated with an increase in cardiovascular deaths in these patients, especially with escalating doses<sup>4</sup> and the late peaks in cortisol might be harmful<sup>11</sup>. A delayed release preparation (Chronocort) taken last thing at night<sup>12</sup> has the theoretical advantage of mimicking the early morning 4am rise in cortisol, which cannot be achieved with other preparations. Chronocort has already shown potential to suppress morning and 24-hour androstenedione levels in patients with congenital adrenal hyperplasia during phase II trials<sup>13</sup>, but more studies are needed to determine the clinical importance of this anticipatory rise in cortisol. **Unfortunately in the most recent trials, Chronocort was unable to achieve its primary outcome of showing superiority over standard care in congenital adrenal hyperplasia during phase III trials, and further development on it is currently on hold.**

Another strategy available in some countries is the use of dual release hydrocortisone (Plenadren) once daily first thing in the morning. This sustained release preparation fails to replicate the pre-awakening rise in cortisol and we know very little about what happens to blood or tissue levels of cortisol overnight. It relies on continuous absorption of hydrocortisone from the gut which can be variable, a problem that must be guarded against in patients with infectious diarrhoea, when glucocorticoids are crucial<sup>14</sup>. The manufacturers state that Plenadren is commonly associated (> 1 in 10 cases) with gastrointestinal side effects<sup>15</sup>.

Building on the early comparative findings that Plenadren may lower blood pressure, weight and improve HbA1c compared to multiple daily cortisone acetate or hydrocortisone<sup>16</sup>, the DREAM study undertook a similar comparison<sup>17</sup>. Although the primary outcome focussed on weight change, the

DREAM study also looked at the effect of Plenadren on immune profiles. Recruiting 89 patients on multiple daily cortisone acetate or hydrocortisone, the investigators switched 46 patients to Plenadren and asked 43 to continue on the thrice daily immediate release preparations. A third group of 25 healthy volunteers were also recruited and data collected over a 24-week follow up period. In addition to a significant corrected weight loss of 4Kg in the switch group, the investigators reported significantly higher classical (CD14+CD16-) monocytes, with lower non-classical (CD14-CD16+) monocytes and CD16+ mature natural killer cells in patients taking thrice-daily hydrocortisone at baseline versus the healthy volunteers. Converting patients to once daily Plenadren was associated with improved numbers of the classical and non-classical monocytes and the mature natural killer cells, with concordant decreases in the soluble CD16 immune marker (an indicator of natural killer cell immaturity) and reported rates of infection as measured by a questionnaire<sup>17</sup>. These findings suggest that Plenadren may be less immunosuppressive than multiple daily dosing of glucocorticoids and that leucocytes may hold the key to assessing the adequacy of an individual's glucocorticoid replacement therapy. Because a majority of the patients were on cortisone acetate, it is not clear whether the variable metabolism of cortisone to cortisol might be a factor.

A DREAM ancillary study also investigated the effect of the switch to Plenadren on the RNA expression of clock genes. The analysis found that 19 of 68 different clock genes were differently expressed at baseline in all patients taking thrice daily hydrocortisone compared to the healthy volunteer control group<sup>18</sup>. In the switch group, the expression of 18 of the 19 affected genes were found to normalised to the levels seen within the healthy volunteers over 12 weeks. Clock genes are involved in auto-regulated feedback cycles thought to be responsible for conveying clock time to individual tissues and cells. The oscillations of clock genes regulate and control the transcription of proteins and can inform tissue sensitivity to glucocorticoids depending on the time of day, through acetylation of lysine residues on the glucocorticoid receptor<sup>6,19</sup>. Although dissociation of peripheral clock genes and the central suprachiasmatic clock is associated with disease states, the functional significance is not understood<sup>19</sup>. The physiological improvements demonstrated by Plenadren have been attributed to its ability to better mirror the natural circadian cortisol profile, compared to multiple daily hydrocortisone or cortisone acetate although this association might simply be due to the lower steroid exposure seen with Plenadren, with data indicating a reduction of approximately 20% in area-under-the-curve exposure<sup>16</sup>. A majority of the patients were on cortisone acetate, which depends on hepatic activation by 11 $\beta$  hydroxysteroid dehydrogenase (11 $\beta$ HSD1) to cortisol<sup>20</sup>. It was assumed that 1mg of cortisone acetate was equivalent to 0.8mg of hydrocortisone<sup>17</sup> but, there is a large (21% to 95%) variability in bioavailability of cortisone acetate<sup>21</sup> as well as tissue specific expression of 11 $\beta$ HSD1.

Prednisolone<sup>22</sup> has a similar plasma profile to Plenadren and since this is taken on waking, the 4am anticipatory rise in glucocorticoid does not occur. At a molecular level we know that prednisolone has prolonged binding to the glucocorticoid receptor compared to hydrocortisone, which would be expected to result in different transcriptional responses in glucocorticoid responsive tissues and prolonged effect<sup>23</sup>. Whether this theoretical advantage is clinically useful and results in reduced risk of adrenal crisis is unknown. The rate of adrenal crises has grown with the increased prescription of short-acting glucocorticoids such as hydrocortisone, although a causal link needs to be confirmed<sup>24</sup>. Further data from a prospective observational study in Germany recorded the frequency of crises experienced by patients on different treatment regimens, and showed that the proportion of patients on prednisolone experiencing a crisis was not greater than expected<sup>25</sup>. It is however clear that previously used doses have been excessive and 3mg to 4mg once daily is sufficient replacement in most patients<sup>26,27</sup>. Currently there are two NIHR adopted studies actively recruiting, comparing low-dose prednisolone with standard regimens of hydrocortisone<sup>28,29</sup>. Low dose prednisolone has been



shown to be superior in reducing androgens, 17-hydroxy progesterone and improving growth velocity in patients with congenital adrenal hyperplasia compared to thrice daily hydrocortisone <sup>30</sup>.

### *Subcutaneous glucocorticoid infusions*

Infusion pumps have been used to mimic the circadian rise in morning cortisol. In addition to normalising morning ACTH levels in patients with primary adrenal failure, improvements in physical functioning and vitality domains of the SF-36 have been noted <sup>31,32</sup>. There are however no further anthropometric or objective benefits, and in double-blind placebo-controlled conditions when compared to oral hydrocortisone treatment, the subjective health outcome benefits are no longer detectable <sup>33</sup>. These studies were performed without any pulsatile or ultradian patterns.

Like many other hormones cortisol is secreted in a pulsatile manner <sup>34,35</sup>. This results in a circadian rhythm that is made up of underlying oscillations which reach maximum amplitude at the time of the circadian peak. In animal models these pulses have been shown to be very important for normal physiological and behavioural activity<sup>36,37</sup>. Although there is relatively little variation of circadian rhythm between subjects, there is much more intra-individual variation in the daily pattern of ultradian pulses<sup>34</sup>. Since the frequency of ultradian pulses simply depends on fixed delays in the feedforward-feedback relationship between the pituitary corticotropes and the adrenal fasciculata cells<sup>38,39</sup> the circadian change in cortisol is determined by changes in pulse amplitude rather than pulse frequency. This provides us with a much easier opportunity to use pumps to replicate physiological pulses, as demonstrated by Russell and colleagues<sup>40</sup>.

It is not yet clear to what extent it is important to replicate these rhythms to provide optimal treatment for patients with adrenal failure. Although we know that stress increases the size of ultradian pulses <sup>41,42</sup>, there have been no long term studies on the modulation of ultradian rhythmicity in normal daily living. Physiological studies have shown that it is the presence of both peaks and troughs of cortisol that is important for optimal regulation of behaviour and of glucocorticoid and mineralocorticoid receptors in animals <sup>43,44</sup>. We are now testing this in new psychological paradigms in man.

From a patient's perspective infusion pumps are less convenient than tablets. However, the technological advances of new miniature nanopumps either external or with subcutaneous reservoirs will make pumps less intrusive. Kalafatakis and colleagues recently undertook a double-blind randomised crossover study, in which they recruited 15 young males and implemented a "block and replace" approach to cortisol using three distinct regimens to investigate the comparative effects of pulsatility on cognition, emotion and sleep <sup>45</sup>. Participants had their endogenous cortisol secretion blocked using metyrapone and received 20mg of hydrocortisone in three modalities: 1- a subcutaneous pulsatile mode to replicate the ultradian profile; 2- a subcutaneous cutaneous infusion; 3- oral thrice daily replacement, each for a five-day period. The order in which the modalities were administered was randomised.

The study demonstrated that pulsatility directly affects neurological physiology in the non-stressed state, independent of the cumulative glucocorticoid exposure. Specifically, subcutaneous ultradian pulsatile infusions promoted better quality of sleep and improved performance of working memory compared to the continuous infusions, as assessed by a sleep questionnaire and functional tests respectively.



These different patterns of cortisol oscillations also had differential effects on the fMRI and functional connectivity of the brain regions which process emotional responses (amygdala, dorsal striatum, insula and orbitofrontal cortex) – with parallel changes detected in attentional bias to, and recognition accuracy of emotional cues<sup>45</sup>.

These data imply that oscillating levels of cortisol are important for physiological regulation of glucocorticoid receptors and post receptor signalling in the brain and for the regulation of mood in man. Whether differences in endogenous patterns of pulsatility in different individuals alters these functions is unknown. The importance of these pulses in longer term replacement therapy in patients with Addison’s disease will be clearer once the Medical Research Council PULSES trial is complete (MR/R010919/1).

Neither immediate release hydrocortisone nor any other oral glucocorticoid will ever replace normal ultradian secretion which remains highly variable across individuals.

**Conclusion**

Long term patient health and cognitive and emotional state as well as patient convenience and compliance are crucial and a balance between a once daily oral dosing, more frequent oral dosing and the requirement of a pump needs to be considered. Currently there are no good studies comparing low dose prednisolone, dual release hydrocortisone (Plenadren, Duocort), delayed release hydrocortisone (Chronocort) and miniature ultradian pumps with three times daily hydrocortisone, using currently understood lower and biologically equivalent doses. We hope that well controlled, relatively long-term double-blind studies will soon be available to allow a rational approach to optimal therapy on an individual patient basis.

**References:**

1. DUNLOP D. Eighty-six cases of addison's disease. *Br Med J*. 1963;2(5362):887-891.
2. Bancos I, Hazeldine J, Chortis V, et al. Primary adrenal insufficiency is associated with impaired natural killer cell function: A potential link to increased mortality. *Eur J Endocrinol*. 2017;176(4):471-480.
3. Behan LA, Carmody D, Rogers B, et al. Low-dose hydrocortisone replacement is associated with improved arterial stiffness index and blood pressure dynamics in severely adrenocorticotrophin-deficient hypopituitary male patients. *Eur J Endocrinol*. 2016;174(6):791-799.
4. Sherlock M, Reulen RC, Alonso AA, et al. ACTH deficiency, higher doses of hydrocortisone replacement, and radiotherapy are independent predictors of mortality in patients with acromegaly. *J Clin Endocrinol Metab*. 2009;94(11):4216-4223.

- 1  
2  
3 5. Hammarstrand C, Ragnarsson O, Hallen T, et al. Higher glucocorticoid replacement doses are  
4 associated with increased mortality in patients with pituitary adenoma. *Eur J Endocrinol*.  
5  
6 2017;177(3):251-256.  
7  
8  
9
- 10  
11 6. Nicolaides NC, Charmandari E, Kino T, Chrousos GP. Stress-related and circadian secretion and  
12 target tissue actions of glucocorticoids: Impact on health. *Front Endocrinol (Lausanne)*. 2017;8:70.  
13  
14  
15
- 16 7. Charmandari E, Nicolaides NC, Chrousos GP. Adrenal insufficiency. *Lancet*. 2014;383(9935):2152-  
17 2167.  
18  
19  
20
- 21 8. Charmandari E, Chrousos GP, Lambrou GI, et al. Peripheral CLOCK regulates target-tissue  
22 glucocorticoid receptor transcriptional activity in a circadian fashion in man. *PLoS One*.  
23  
24 2011;6(9):e25612.  
25  
26  
27
- 28 9. Cuesta M, Cermakian N, Boivin DB. Glucocorticoids entrain molecular clock components in human  
29 peripheral cells. *FASEB J*. 2015;29(4):1360-1370.  
30  
31  
32
- 33 10. Monk TH, Buysse DJ. Exposure to shift work as a risk factor for diabetes. *J Biol Rhythms*.  
34  
35 2013;28(5):356-359.  
36  
37  
38
- 39 11. Muller L, Quinkler M. Adrenal disease: Imitating the cortisol profile improves the immune  
40 system. *Nat Rev Endocrinol*. 2018;14(3):137-139.  
41  
42  
43
- 44 12. Whitaker M, Debono M, Huatan H, Merke D, Arlt W, Ross RJ. An oral multiparticulate, modified-  
45 release, hydrocortisone replacement therapy that provides physiological cortisol exposure. *Clin*  
46  
47 *Endocrinol (Oxf)*. 2014;80(4):554-561.  
48  
49  
50
- 51 13. Mallappa A, Sinaii N, Kumar P, et al. A phase 2 study of chronocort, a modified-release  
52 formulation of hydrocortisone, in the treatment of adults with classic congenital adrenal hyperplasia.  
53  
54  
55  
56  
57  
58  
59  
60 *J Clin Endocrinol Metab*. 2015;100(3):1137-1145.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

14. Allolio B. Extensive expertise in endocrinology. adrenal crisis. *Eur J Endocrinol.* 2015;172(3):R115-24.

15. Shire Pharmaceuticals. Summary of product characteristics.  
[https://www.ema.europa.eu/documents/product-information/plenadren-epar-product-information\\_en.pdf](https://www.ema.europa.eu/documents/product-information/plenadren-epar-product-information_en.pdf).

16. Johannsson G, Nilsson AG, Bergthorsdottir R, et al. Improved cortisol exposure-time profile and outcome in patients with adrenal insufficiency: A prospective randomized trial of a novel hydrocortisone dual-release formulation. *J Clin Endocrinol Metab.* 2012;97(2):473-481.

17. Isidori AM, Venneri MA, Graziadio C, et al. Effect of once-daily, modified-release hydrocortisone versus standard glucocorticoid therapy on metabolism and innate immunity in patients with adrenal insufficiency (DREAM): A single-blind, randomised controlled trial. *Lancet Diabetes Endocrinol.* 2017.

18. Venneri MA, Hasenmajer V, Fiore D, et al. Circadian rhythm of glucocorticoid administration entrains clock genes in immune cells: A DREAM trial ancillary study. *J Clin Endocrinol Metab.* 2018;103(8):2998-3009.

19. Oster H, Challet E, Ott V, et al. The functional and clinical significance of the 24-hour rhythm of circulating glucocorticoids. *Endocr Rev.* 2017;38(1):3-45.

20. Fariss BL, Hane S, Shinsako J, Forsham PH. Comparison of absorption of cortisone acetate and hydrocortisone hemisuccinate. *J Clin Endocrinol Metab.* 1978;47(5):1137-1140.

21. Heazelwood VJ, Galligan JP, Cannell GR, Bochner F, Mortimer RH. Plasma cortisol delivery from oral cortisol and cortisone acetate: Relative bioavailability. *Br J Clin Pharmacol.* 1984;17(1):55-59.

22. Williams EL, Choudhury S, Tan T, Meeran K. Prednisolone replacement therapy mimics the circadian rhythm more closely than other glucocorticoids. *The Journal of Applied Laboratory Medicine*. 2016;1(2):152-161.
23. Stavreva DA, Wiench M, John S, et al. Ultradian hormone stimulation induces glucocorticoid receptor-mediated pulses of gene transcription. *Nat Cell Biol*. 2009;11(9):1093-1102.
24. Rushworth RL, Torpy DJ. Adrenal insufficiency in australia: Is it possible that the use of lower dose, short-acting glucocorticoids has increased the risk of adrenal crises? *Horm Metab Res*. 2015;47(6):427-432.
25. Hahner S, Spinnler C, Fassnacht M, et al. High incidence of adrenal crisis in educated patients with chronic adrenal insufficiency: A prospective study. *J Clin Endocrinol Metab*. 2015;100(2):407-416.
26. Smith DJF, Prabhudev H, Choudhury S, Meeran K. Prednisolone has the same cardiovascular risk profile as hydrocortisone in glucocorticoid replacement. *Endocr Connect*. 2017;6(8):766-772.
27. Choudhury S, Machenahalli P, Tan T, Meeran K. Inadvertent treatment of hypoadrenalism with prednisolone in pemphigus: A case report. *Clin Case Rep*. 2019;0(0).
28. Imperial College London. Safety and efficacy of prednisolone in adrenal insufficiency disease (PRED-AID study). ISRCTN Registry Web site. <http://www.isrctn.com/ISRCTN41325341>. Published 01/28/2019. Updated 2019. Accessed 04/19, 2019.
29. Imperial College London. **Hydrocortisone vs prednisolone in AI (HYPER-AID)**. Clinicaltrials.gov Web site. <https://clinicaltrials.gov/ct2/show/NCT03608943>. Published 08/01/2018. Updated 2019. Accessed 04/19, 2019.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

30. Caldato MC, Fernandes VT, Kater CE. One-year clinical evaluation of single morning dose prednisolone therapy for 21-hydroxylase deficiency. *Arq Bras Endocrinol Metabol.* 2004;48(5):705-712.

31. Lovas K, Husebye ES. Continuous subcutaneous hydrocortisone infusion in addison's disease. *Eur J Endocrinol.* 2007;157(1):109-112.

32. Oksnes M, Bjornsdottir S, Isaksson M, et al. Continuous subcutaneous hydrocortisone infusion versus oral hydrocortisone replacement for treatment of addison's disease: A randomized clinical trial. *J Clin Endocrinol Metab.* 2014;99(5):1665-1674.

33. Gagliardi L, Nenke MA, Thynne TR, et al. Continuous subcutaneous hydrocortisone infusion therapy in addison's disease: A randomized, placebo-controlled clinical trial. *J Clin Endocrinol Metab.* 2014;99(11):4149-4157.

34. Lightman S, Terry JR. The importance of dynamic signalling for endocrine regulation and drug development: Relevance for glucocorticoid hormones. *Lancet Diabetes Endocrinol.* 2014;2(7):593-599.

35. Faghih RT, Dahleh MA, Adler GK, Klerman EB, Brown EN. Deconvolution of serum cortisol levels by using compressed sensing. *PLoS One.* 2014;9(1):e85204.

36. Lightman SL, Conway-Campbell BL. The crucial role of pulsatile activity of the HPA axis for continuous dynamic equilibration. *Nat Rev Neurosci.* 2010;11(10):710-718.

37. Spiga F, Walker JJ, Terry JR, Lightman SL. HPA axis-rhythms. *Compr Physiol.* 2014;4(3):1273-1298.

38. Walker JJ, Terry JR, Lightman SL. Origin of ultradian pulsatility in the hypothalamic-pituitary-adrenal axis. *Proc Biol Sci.* 2010;277(1688):1627-1633.

39. Walker JJ, Spiga F, Waite E, et al. The origin of glucocorticoid hormone oscillations. *PLoS Biol.* 2012;10(6):e1001341.
40. Russell GM, Durant C, Ataya A, et al. Subcutaneous pulsatile glucocorticoid replacement therapy. *Clin Endocrinol (Oxf).* 2014;81(2):289-293.
41. Henley DE, Russell GM, Douthwaite JA, et al. Hypothalamic-pituitary-adrenal axis activation in obstructive sleep apnea: The effect of continuous positive airway pressure therapy. *J Clin Endocrinol Metab.* 2009;94(11):4234-4242.
42. Gibbison B, Spiga F, Walker JJ, et al. Dynamic pituitary-adrenal interactions in response to cardiac surgery. *Crit Care Med.* 2015;43(4):791-800.
43. Sarabdjitsingh RA, Conway-Campbell BL, Leggett JD, et al. Stress responsiveness varies over the ultradian glucocorticoid cycle in a brain-region-specific manner. *Endocrinology.* 2010;151(11):5369-5379.
44. George CL, Birnie MT, Flynn BP, Kershaw YM, Lightman SL, Conway-Campbell BL. Ultradian glucocorticoid exposure directs gene-dependent and tissue-specific mRNA expression patterns in vivo. *Mol Cell Endocrinol.* 2017;439:46-53.
45. Kalafatakis K, Russell GM, Harmer CJ, et al. Ultradian rhythmicity of plasma cortisol is necessary for normal emotional and cognitive responses in man. *Proc Natl Acad Sci U S A.* 2018;115(17):E4091-E4100.